

Substituting Values Into Equations

When substituting values into equations, make sure to use the correct order of operations:

- brackets first
- multiply and divide in order from left to right
- add and subtract in order from left to right

Determine the value of y when substituting $x = 7$ into the following equation.

$$y = 2(x - 3) + 5$$

$$y = 2(7 - 3) + 5 \quad \text{Brackets}$$

$$y = 2(4) + 5 \quad \text{Multiply}$$

$$y = 8 + 5 \quad \text{Add}$$

$$y = 13$$

1. Determine the value of y in each equation when $d = 6$.
 - a) $y = (3d + 4) \div 2 + 8$
 - b) $y = 5 \times 4 + d \div 2$
 - c) $y = (3 + d - 7) \times 4d + 5$
2. Calculate the surface area of each rectangular prism using the formula $SA = 2(bh + bl + hl)$, where b = base, h = height, and l = length.
 - a) $b = 5$ cm, $h = 11$ cm, $l = 12$ cm
 - b) $b = 6$ cm, $h = 10$ cm, $l = 9$ cm
 - c) $b = 7$ cm, $h = 12$ cm, $l = 6$ cm

Modelling and Solving One-Step Equations

To solve a problem, you sometimes need to translate words into equations. For example, "the *sum* of 4 and another number is 12" can be modelled by the equation $4 + x = 12$.

The equation can now be solved.

$$x + 4 = 12$$

$$x + 4 - 4 = 12 - 4 \quad \text{Subtract 4 from both sides of the equation.}$$

$$x = 8$$

3. Model each situation with an algebraic equation.
 - a) seven more than a number, p , is twelve
 - b) three less than a number, x , is eleven
 - c) four times a number, s , is twenty-eight
 - d) when a number, k , is divided by six, the result is nine

4. Develop and solve an algebraic equation for each question.
- a) You share \$20 equally among four people. How much does each person get?
- b) If Jim's height increased by 13 cm over the past year and he is now 152 cm, how tall was he a year ago?
- c) After Lynn triples the amount she has in her bank account, she has \$54. How much did she initially have in her account?
- d) Ayisha worked twice as long on a math project as Harpreet did. If Ayisha worked for 50 min on the project, how long did Harpreet work on it?

Solving Two-Step Equations

To solve a two-step problem of the form $ax + b = c$, you need to isolate the variable on one side of the equal sign. When undoing the operations performed on the variable, follow the reverse order of operations:

- Subtract and add in order from left to right.
- Multiply and divide in order from left to right.

Solve $6x + 7 = 25$.

$$6x + 7 = 25$$

$$6x + 7 - 7 = 25 - 7 \quad \text{Subtract 7 from both sides of the equation.}$$

$$6x = 18$$

$$\frac{6x}{6} = \frac{18}{6}$$

Divide both sides of the equation by 6.

$$x = 3$$

Check:

$$\text{Left Side} = 6x + 7 \quad \text{Right Side} = 25$$

$$= 6(3) + 7$$

$$= 18 + 7$$

$$= 25$$

$$\text{Left Side} = \text{Right Side}$$

The solution is correct.

5. Copy each equation. Circle the first operation you undo. Underline the second operation you undo.
- a) $2n + 4 = 18$ b) $3x + 5 = 17$
- c) $8y - 70 = 94$ d) $27 = 7q + 6$
6. Solve each equation. Check your solution.
- a) $9 + 5j = 49$ b) $4t + 2 = 14$
- c) $10x - 7 = 23$ d) $6p - 5 = 31$
7. Jan is sewing leather trim and ribbon on a new parka. The length of the leather trim is 40 cm more than 4 times the length of the ribbon. If Jan uses 240 cm of trim, how much ribbon does she use?